

GI Bleeding – Upper Admission

(orders in addition to General Admission orders)

Vital Signs & Interventions

- Measure blood pressure, orthostatic
- Communication order lie patient on side
- Nasogastric/orogastric tube insertion/management
- Oral suction at bedside

Medications

Antacids

- calcium carbonate 500 mg PO 4 times a day as needed for dyspepsia
- MAALOX PLUS ES 15 mL PO every 4 hours as needed for dyspepsia
- MYLANTA 30 mL PO every 4 hours as needed for dyspepsia

Histamine-2 Receptor Antagonists

- Famotidine (PEDCID) 20 mg IV every 12 hours 20 mg PO 2 times a day

Proton Pump Inhibitors

- esomeprazole (NEXIUM) 40 mg IV daily
- omeprazole (PRILOSEC) 20 mg PO daily

Beta-Blockers

- atenolol (TENORMIN) 50 mg PO daily
- Propranolol (INDERAL) 40 mg PO 2 times a day 80 mg PO 2 times a day
- propranolol SR 80 mg 1 capsule PO daily

Laboratory

Blood Bank

- Type & crossmatch, immediate spin Type & screen Transfusion, red blood cells

Hematology

- Occult blood, stool

Serology

- Helicobacter pylori antibody

Other

- Urea breath test

Diagnostic Tests

Computed Tomography

- CT, abdomen, with contrast
- CT, abdomen, without contrast
- CT, chest, with contrast
- CT, chest, without contrast

General Radiography

- Radiograph, upper GI series

Nuclear Medicine

- Nuclear medicine, gastrointestinal blood loss imaging

Reminders: Consider elective intubation for patients with acute variceal bleeding or patients who are actively vomiting

Consults

- Consult gastroenterology Dr. _____
- Consult general surgery Dr. _____
- Consult interventional radiology Dr. _____
- Consult to social services

Provider Signature _____ **Date and Time** _____

This order set is not built in FAMIS

Patient Information

St. Elizabeth's Hospital
Belleville, Illinois
GI Bleed-Upper Admission

Form # 1409

Org. 01/09, Rev. 6/09,5/10

Page 1 of 1

