


**GI (Biliary Tract) Pre-operative - DEBMPRBILI**

1.  orders are active unless order is manually lined through to inactivate
2.  do not enter as order unless checked
3. • **REMINDER:** do not enter as order

**Interventions**

- Confirm signed consent on chart for \_\_\_\_\_
- Notify medical records to send results to primary care physician
- Elastic stockings, below the knee
- Sequential compression pneumatic devices (SCDs) / Pulsatile Anti-Embolism System (PAS) - discontinue PAS when patient is ambulating without assistance 

**Patient/Caregiver Education**

- Education: deep breathing & coughing exercises
- Education: incentive spirometry
- Education: Wound care

**Diet**

- NPO after midnight on \_\_\_\_/\_\_\_\_/\_\_\_\_
- Other \_\_\_\_\_

**Laboratory** (Indicate schedule following order)

- Type & screen for surgery today
- Complete blood cell count with automated white blood cell differential
- Comprehensive metabolic panel (CMP - CHEM 14)
- Pregnancy urine, if female is of childbearing age & has pregnancy ability
- Basic metabolic panel (BMP - CHEM 8) \_\_\_\_\_
- Prothrombin time (PT) & international normalized ratio (INR) & Partial thromboplastin time (PTT), if on Coumadin
- Amylase, serum \_\_\_\_\_
- Lipase \_\_\_\_\_
- Urinalysis (UA) with microscopy \_\_\_\_\_
- Urinalysis (UA) with reflex to culture \_\_\_\_\_
- Other \_\_\_\_\_

**Diagnostic Tests (If not requested by anesthesia. Check with Anesthesia)**

- Chest X-Ray, PA & Lateral
  - Now (indication) \_\_\_\_\_
  - in AM (indication) \_\_\_\_\_
- 12-lead ECG
- Other: \_\_\_\_\_

**Medications**

- VERIFY HOME MEDICATIONS

**Antibiotics (IV)**

- **Pre-op antibiotic should be given in the Operating Room WITHIN ONE HOUR of surgical incision**
  - cefazolin (ANCEF) 1 gram IV x1 dose

**Antibiotics (B-lactam Allergy)-IV**

- levofloxacin (LEVAQUIN) 500 mg IV x1 dose  
**\*\*AND\*\***  
clindamycin (CLEOCIN) 600 mg IV x1 dose

**Beta-Blockers** 

- Patients undergoing noncardiac surgery who are currently on beta-blocker therapy &/or who are at high risk for coronary artery disease should receive a beta-blocker during the perioperative period
  - metoprolol 5 mg IV q6h PRN to achieve target heart rate of \_\_\_\_\_

**Provider Signature**

\_\_\_\_\_

**Date** \_\_\_\_\_ **Time** \_\_\_\_\_

 = Performance Measure

Patient Information

St. Elizabeth's Hospital  
Belleville, Illinois

**GI-Biliary Tract Pre-op**

Form # 1455

Org. 7/09 Rev. 8/10

Page 1 of 1



N.04SU